

EFFECTS OF FIRE ON GOLDEN EAGLE TERRITORY OCCUPANCY AND REPRODUCTIVE SUCCESS

MICHAEL N. KOCHERT,¹ U.S. Geological Survey, Forest and Rangeland Ecosystem Science Center, Snake River Field Station, 970 Lusk, Boise, ID 83706, USA

KAREN STEENHOF, U.S. Geological Survey, Forest and Rangeland Ecosystem Science Center, Snake River Field Station, 970 Lusk, Boise, ID 83706, USA

LESLIE B. CARPENTER, Raptor Research Center, Boise State University, Boise, ID 83725, USA

JOHN M. MARZLUFF,² Greenfalk Consultants, 8210 Gantz, Boise, ID 83709, USA

Abstract: We examined effects of fire on golden eagle (*Aquila chrysaetos*) territory occupancy and reproductive success in southwestern Idaho because wildfires since 1980 have resulted in large-scale losses of shrub habitat in the Snake River Plain. Success (percentage of pairs that raised young) at burned territories declined after major fires ($P = 0.004$). Pairs in burned areas that could expand into adjacent vacant territories were as successful as pairs in unburned territories and more successful than pairs in burned territories that could not expand. Success at extensively burned territories was lowest 4–6 years after burning but increased 4–5 years later. The incidence and extent of fires did not help predict territories that would have low occupancy and success rates in postburn years. The presence of a vacant neighboring territory and the amount of agriculture and proportion of shrubs within 3 km of the nesting centroid best predicted probability of territory occupancy. Nesting success during preburn years best predicted the probability of a territory being successful in postburn years. Burned territories with high success rates during preburn years continued to have high success rates during postburn years, and those with low success in preburn years continued to be less successful after burning. In areas where much shrub habitat has been lost to fire, management for golden eagles should include active fire suppression and rehabilitation of burned areas.

JOURNAL OF WILDLIFE MANAGEMENT 63(3):773–780

Key words: *Aquila chrysaetos*, fire, golden eagle, habitat selection, home range, Idaho, reproductive success, shrub-steppe, territory occupancy.

Most raptor studies have focused on immediate or short-term responses of wintering or foraging raptors to fire (Komarek 1969, Smallwood et al. 1982, Tewes 1984, Chavez-Ramirez and Prieto 1994); long-term effects of fire on nesting raptors are poorly understood. Watson et al. (1987) predicted that burning may reduce golden eagle prey by eliminating shrub habitat. Predictions about the effects of fire or other habitat alteration on breeding raptor populations are complicated by the interaction between habitat quality and attributes of the pair (Newton 1991, 1992). Nesting success depends not only on habitat but also on the individuals that live there.

Golden eagles typically use and defend a group of 1–13 nests and a surrounding hunting range collectively known as a “territory” (Watson 1997). Golden eagles occupy territories year-round in southwestern Idaho, and the same individuals tend to occupy the same ter-

ritory year after year (Marzluff et al. 1997, Watson 1997). Golden eagles nesting in southwestern Idaho prey primarily on black-tailed jackrabbits (*Lepus californicus*; Steenhof and Kochert 1988), and golden eagle reproduction is closely tied to jackrabbit abundance (Steenhof et al. 1997). Black-tailed jackrabbits in southwestern Idaho are associated with shrub habitats, and jackrabbit densities are low in disturbed grassland habitats (Smith and Nydegger 1985, Knick and Dyer 1997). In the early 1980s, wildfires in southwestern Idaho burned vast areas of shrubs, resulting in large stands of exotic annual grasses (primarily cheatgrass [*Bromus tectorum*]) and forbs. The reduction in black-tailed jackrabbit habitat within the foraging areas of nesting golden eagles provided a unique opportunity to study long-term effects of fire on occupancy and nesting success of golden eagles.

In this paper, we examine 24 years of data on golden eagle territory occupancy and reproductive success in relation to the occurrence of wildfires. Because of the relation between black-tailed jackrabbits and shrub habitats, and the dependence of golden eagles on jackrabbits,

¹ E-mail: mkochert@eagle.idbsu.edu

² Present address: College of Forest Resources, University of Washington, Seattle, WA 98195, USA.

layers depicted burns, 1979 and 1994 vegetation types, and agriculture. The burn layer was digitized from 1:24,000-scale U.S. Geological Survey topographic maps depicting boundaries of fires that occurred from 1950 to 1994 (U.S. Department of the Interior 1996). The 1979 vegetation map was developed via visual interpretation of low-level aerial photographs and field verification of vegetation stands (U.S. Department of the Interior 1979), and the 1994 vegetation map was developed via supervised classification of Landsat thematic mapper satellite imagery (Knick *et al.* 1997). The agricultural layer depicted all agricultural lands, including fallow fields, and was a composite of the 1979 and 1994 vegetation layers and 1993 U.S. Bureau of Reclamation agricultural maps (U.S. Department of the Interior 1996).

We computed the area of each habitat and disturbance characteristic within each proximity ring. We calculated the proportion of area that burned each year and the total proportion burned since 1950, excluding areas that reburned. We used the 1994 NCA vegetation map to determine the percentage of area covered by shrubs in postburn years.

Statistical Analyses

Because specific pre- and postburn years varied by territory and annual success rates of golden eagles vary significantly with prey and weather conditions (Steenhof *et al.* 1997), we adjusted for "year effect" in pre- and postburn analyses. We evaluated occupancy and success of each burned territory during years before and after it burned, relative to the 9 unburned territories in the same corresponding years. We computed standardized scores for each territory during both the preburn and postburn periods. Standardized scores were the difference between mean occupancy or success for the burned territory during the particular time period and mean occupancy or success for the 9 unburned territories during the same years. For example, mean preburn nesting success for the Commeford territory from 1971 to 1983 was 75.0%, while mean success for all unburned territories during the same years was 50.5%. Thus the preburn score was 24.5 (75.0–50.5). Mean success for postburn years (1984–94) was 54.5% for the Commeford territory and 44.9% for all unburned territories, and the postburn score was 9.6 (54.5–44.9).

We used Wilcoxon signed rank tests (Z) to

determine whether standardized occupancy or success scores decreased after the territory burned. We used stepwise logistic regression to identify variables important in predicting occupancy and success of golden eagle territories in postburn years. We included 3 habitat variables (percentage of the area in shrubs, in agriculture, and burned as of 1994) and 2 golden eagle variables (mean 1971–81 nesting success and whether the nearest adjacent territory was vacant during postburn years) in stepwise logistic regression models predicting occupancy and success from 1986 to 1994.

We used SYSTAT (Wilkinson *et al.* 1994, SPSS 1997) and StatXact (Cytel Software 1989) for statistical analyses and considered P -values ≤ 0.05 as significant. Because we predicted the direction of changes in occupancy and success in relation to the habitat and disturbance variables, we used 1-tailed tests unless otherwise stated.

RESULTS

Fire Occurrence

From 1950 to 1994, fires burned 0–100% (\bar{x} = 35%, median = 29) of the area within the proximity rings of 36 historical golden eagle territories in the NCA. We used the mean proportion of area burned to distinguish between extensively and moderately burned territories and classified 17 as extensively burned (\bar{x} = 61%, range = 36–100%), 10 as moderately burned (\bar{x} = 22%, range = 11–29%), and 9 as unburned (\bar{x} = 0.3%, range = 0–2%). The incidence of fires was relatively low prior to 1980. Only 14 territories burned between 1950 and 1979. These 14 territories and 13 new ones burned during the 1980s; all but 1 burned between 1981 and 1985. Fires in the 1980s were larger and more numerous than in the 1970s; a mean of 50% of the area within 3 km of nesting centroids burned in the 1980s compared to 7% during the 1970s. From 1990 to 1994, territories essentially remained unburned, with $\leq 3.3\%$ of the area burned around only 5 nesting centroids.

Postfire Changes in Occupancy and Success

Mean standardized occupancy scores for all 27 burned territories did not change after burning (Table 1; Z = -0.38 , P = 0.36); however, effect of burning on occupancy scores was related to whether or not pairs could expand into

Table 2. Results of 2 separate logistic regression procedures predicting occupancy and success at golden eagle territories in postburn years. The first model distinguishes territories occupied in all years ($n = 26$) from those vacant at least 1 year ($n = 10$), 1986–94. The second model distinguishes territories successful in >5 years ($n = 14$) from those successful <5 years ($n = 21$), 1986–94^a. One territory was excluded from the success model due to incomplete data.

Variable	Parameter estimate ^b	SE	<i>t</i> -ratio	<i>P</i> -value	Odds ratio
Occupancy					
Intercept	-2.647	1.590	-1.665	0.096	
% shrub	0.081	0.032	2.540	0.011	1.085
% agriculture	0.107	0.050	2.154	0.031	1.113
Vacant neighbor	1.710	0.786	2.177	0.030	5.529 ^c
Success					
Intercept	-2.933	1.165	-2.517	0.012	
1971–81 success	0.049	0.021	2.382	0.017	1.051

^a Variables shown are those that contributed significantly to stepwise logistic regression models ($P < 0.05$; SYSTAT, logit procedure). Variables that did not contribute significantly to either model were percentage of area burned and whether or not burns occurred within 3 km of the nesting centroid. Overall significance was 0.04 for the occupancy model ($G_3 = 13.33$) and 0.07 for the success model ($G_1 = 7.15$).

^b Positive values indicate that an increase in the independent variable is associated with a higher probability of a territory being occupied or successful.

^c Territories were 5.5 times more likely to be occupied in postburn years if the neighboring territory was vacant.

Annual nesting success of golden eagles in the NCA ranged from 32 to 80% between 1986 and 1994. Twenty-one territories were successful in $<50\%$ of postburn years. Habitat measures failed to predict which territories would be successful in $>50\%$ of postburn years. The only variable that contributed significantly to a logistic regression model (Table 2) predicting success in postburn years was a territory's nesting success in preburn years (1971–81). Preburn nesting success was positively associated with nesting success in postburn years, and the model was significant ($P = 0.007$).

DISCUSSION

Overall, golden eagles in the NCA responded negatively to the loss of shrubs (black-tailed jackrabbit habitat) that resulted from fires. However, the incidence and extent of fires did not help us to predict which territories would have low occupancy and success rates in postburn years. Responses to fire were variable and influenced by at least 3 factors: (1) whether the nearest neighboring territory was vacant; (2) the ability to use alternative foraging habitat (i.e., farmland, cliff, talus, riparian); and (3) the underlying quality of the pair or territory.

Territories were more likely to be occupied during postburn years if the nearest neighboring territory was vacant, and success of pairs that had the opportunity to extend their range into a neighboring vacant territory did not change after burning. Golden eagle pairs responded to reduced shrub habitat from burning by ranging over larger areas and concentrating

their use in the shrub patches within these larger ranges (Marzluff et al. 1997). Postburn success rates might have been even lower if neighboring territories had not become vacant after burning.

Golden eagles also may have compensated for loss of shrub habitat due to burning by using alternative habitats. Marzluff et al. (1997) observed that some golden eagles in areas with little shrub habitat as a result of burning in the NCA restricted their foraging to small areas of cliff and riparian habitat around their nests. They preyed less on black-tailed jackrabbits and took more alternate prey such as rock doves (*Columba livia*), waterfowl, yellow-bellied marmots (*Marmota flaviventris*), and desert cottontails (*Sylvilagus nuttalli*). Territory occupancy during the postburn years was predicted, in part, by the amount of agriculture within the proximity rings. The positive relation with agriculture could be explained by availability of alternate prey in farmlands. Golden eagles nesting in agricultural areas in southwestern Idaho took more ring-necked pheasants (*Phasianus colchicus*), yellow-bellied marmots, and other prey than golden eagles in the rangelands (Kochert 1972).

Postburn nesting success at individual territories was related primarily to nest success in preburn years. Territories with high success rates during preburn years continued to have high success rates in postburn years. Territory occupancy and breeding success for many bird species are related to quality of the territory (habitat) and of the pair occupying the territory

ations, particularly if cliff, talus, riparian, and farmland habitats are available. However, overall losses to golden eagles will be significant.

Managers need to monitor long-term effects of habitat alteration and the relative values of different habitat patches. Monitoring must include more than a simple count of the number of occupied golden eagle territories because, as populations decline, productivity per pair may actually increase as marginal habitats are no longer used (Ferrer and Donazar 1996). Number of young golden eagles produced per pair did not decline in the NCA from 1971 to 1994, despite a decline in the number of occupied nesting territories and a possible reduced carrying capacity for golden eagles (Steenhof et al. 1997). Additional monitoring is needed to determine if the ability of golden eagles to withstand habitat alteration is a long-term response. Biologists will particularly need to know if some vacant territories will be reoccupied in future years or if more territories will become permanently vacant, and if remaining pairs compensate via increased productivity.

ACKNOWLEDGMENTS

This paper is a contribution of the Snake River Field Station, Forest and Rangeland Ecosystem Science Center, U.S. Geological Survey (formerly the Raptor Research and Technical Assistance Center, U.S. Bureau of Land Management [BLM] and subsequently the National Biological Service [NBS]). The Idaho Cooperative Wildlife Research Unit provided the 1971 golden eagle nesting data. Data from 1972 to 1989 were collected as part of the BLM's Snake River Birds of Prey Research Project. The BLM provided funds from 1972 to 1994. From 1984 to 1989, Pacific Power and Light Company funded the aerial surveys. As part of the Cooperative BLM-Idaho Army National Guard (IDARNG) Research Project, the IDARNG provided funding from 1990 to 1994 under U.S. Army contract D05-90-0135 and numerous agreements administered by W. S. Seegar. Additional funding and support was provided by NBS. J. H. Doremus and A. R. Bammann supervised collection of nesting data from 1976 to 1983, and J. H. Doremus assisted with data collection from 1983 to 1990. R. N. Lehman, M. S. Vekasy, and their crews helped to collect data from 1990 to 1994. The study could not have been completed without the assistance of numerous field technicians and dedicated volun-

teers, especially J. R. Oakley, T. E. Hamer, O. R. Fremming, and M. W. Collopy. Special thanks go to G. R. Griggs, D. W. Dyer, and E. E. Black for logistical support and coordination and to the Snake River Basin Office, U.S. Fish and Wildlife Service, for use of a jet boat. G. W. Smith, N. C. Nydegger, and D. L. Quinney developed the 1979 vegetation map, and S. T. Knick and S. E. Watts supervised collection of the 1994 vegetation data. D. L. Dyer and T. J. Zarriello conducted the GIS analyses and prepared habitat maps, and D. M. Parrish tabulated most of the nesting data. J. T. Rotenberry provided valuable ideas for developing standardized scores for the golden eagle data, and M. W. Collopy, S. T. Knick, B. A. Maurer, M. J. McGrady, J. T. Rotenberry, R. S. Lutz, and 2 anonymous reviewers made valuable comments on earlier drafts of the manuscript.

LITERATURE CITED

- BEECHAM, J. J., JR. 1970. Nesting ecology of the golden eagle in southwestern Idaho. Thesis, University of Idaho, Moscow, Idaho, USA.
- CHAVEZ-RAMIREZ, F., AND F. G. PRIETO. 1994. Effects of prescribed fires on habitat use by wintering raptors on a Texas barrier island grassland. *Journal of Raptor Research* 28:262-265.
- CYTEL SOFTWARE. 1989. StatXact statistical software for exact nonparametric inference. User manual. Cytel Software, Cambridge, Massachusetts, USA.
- FERRER, M., AND J. A. DONAZAR. 1996. Density-dependent fecundity by habitat heterogeneity in an increasing population of Spanish imperial eagles. *Ecology* 77:69-74.
- KNICK, S., AND D. L. DYER. 1997. Distribution of black-tailed jackrabbit habitat determined by GIS in southwestern Idaho. *Journal of Wildlife Management* 61:75-86.
- , J. T. ROTENBERRY, AND T. J. ZARRIELLO. 1997. Supervised classification of Landsat thematic mapper imagery in a semi-arid rangeland by nonparametric discriminant analysis. *Photogrammetric Engineering and Remote Sensing* 63:79-86.
- KOCHERT, M. N. 1972. Population status and chemical contamination of golden eagles in southwestern Idaho. Thesis, University of Idaho, Moscow, Idaho, USA.
- , AND M. PELLANT. 1986. Multiple use in the Snake River Birds of Prey Area. *Rangelands* 8:217-220.
- KOMAREK, E. V. 1969. Fire and animal behavior. *Proceedings of the Tall Timbers Fire Ecology Conference* 9:161-207.
- MARZLUFF, J. M., S. T. KNICK, M. S. VEKASY, L. S. SCHUECK, AND T. J. ZARRIELLO. 1997. Spatial use and habitat selection of golden eagles in southwestern Idaho. *Auk* 114: 673-687.
- , G. E. WOOLFENDEN, J. W. FITZPATRICK, AND R. P. BALDA. 1996. Breeding partnerships of